

original

where  $k = \sqrt{N}$ , and  $N$  is the number of tiles in the system. This free energy, seen in Fig. 8, will increase to a maximum dependent upon  $\varepsilon$  and  $G_{se}$ , creating a barrier to nucleation. As  $\varepsilon$  decreases or  $G_{se}$  increases, the height of the barrier, and the number of tile attachments required to reach it, will increase, with the maximum being reached at  $N = G_{se}^2 / \varepsilon^2$ . At that point, the free energy will be  $G(A) = G_{se} / \varepsilon$ . From eqn (11), the concentration of the critical nucleus will be  $[A]_{crit} = u_0 e^{-G_{se} \varepsilon^{-1} + \alpha}$ . The rate of formation of larger assemblies will be limited by this critical nucleus concentration.

corrected

where  $k = \sqrt{N}$ , and  $N$  is the number of tiles in the system. This free energy, seen in Fig. 8, will increase to a maximum dependent upon  $\varepsilon$  and  $G_{se}$ , creating a barrier to nucleation. As  $\varepsilon$  decreases or  $G_{se}$  increases, the height of the barrier, and the number of tile attachments required to reach it, will increase, with the maximum being reached at  $N = G_{se}^2 / \varepsilon^2$ . At that point, the free energy will be  $G(A) = G_{se}^2 / \varepsilon$ . From eqn (11), the concentration of the critical nucleus will be  $[A]_{crit} = u_0 e^{-G_{se}^2 \varepsilon^{-1} + \alpha}$ . The rate of formation of larger assemblies will be limited by this critical nucleus concentration.